

FLOWERING NEWSLETTER REVIEW

Regulation of flowering time by the miR156-mediated age pathway

Jia-Wei Wang*

National Key Laboratory of Plant Molecular Genetics (NKLPMG), Institute of Plant Physiology and Ecology (SIPPE), Shanghai Institutes for Biological Sciences (SIBS), Shanghai 200032, P. R. China

* To whom correspondence should be addressed. E-mail: jwwang@sibs.ac.cn

Received 25 February 2014; Revised 24 April 2014; Accepted 6 May 2014

Abstract

Precise flowering time is critical to reproductive success. In response to diverse exogenous and endogenous cues including age, hormones, photoperiod, and temperature, the floral transition is controlled by a complex regulatory network, which involves extensive crosstalks, feedback, or feedforward loops between the components within flowering time pathways. The newly identified age pathway, which is controlled by microRNA156 (miR156) and its target *SQUAMOSA PROMOTER BINDING-LIKE (SPL)* transcription factors, ensures plants flower under non-inductive conditions. In this review, I summarize the recent advance in understanding of the age pathway, focusing on the regulatory basis of the developmental decline in miR156 level by age and the molecular mechanism by which the age pathway is integrated into other flowering time pathways.

Key words: Age, microRNA, flowering time.

Introduction

The aerial lateral organs of a plant are derived from the shoot apical meristem (SAM), a population of pluripotent stem cells at the shoot apex that are formed during embryonic development. After seed germination, organ primordia are continuously formed on the flanks of the SAM. Based on the identity of the lateral organs, the post-embryonic development of a plant can be divided into vegetative and reproductive phases. The SAM produces leaves during the vegetative phase, whereas it gives rise to flowers in the reproductive phase (Poethig, 2003). Vegetative phase can be further divided into juvenile and adult phases. Adult phase differs from juvenile phase in terms of reproductive competence and morphological differences such as leaf epidermal cell differentiation and leaf complexity (Huijser and Schmid, 2011; Poethig, 2013).

The floral transition, namely the switch from vegetative to reproductive phase, is coordinately controlled by multiple genetic pathways in response to various developmental and environmental cues (reviewed in Andres and Coupland, 2012; Bäurle and Dean, 2006; Srikanth and Schmid, 2011). The past two decades have seen fundamental advances in our

understanding of the molecular mechanism underlying floral transition. Studies of the annual model *Arabidopsis thaliana* identified five flowering time pathways, known as age, autonomous, gibberellin (GA), photoperiod, and vernalization (Amasino and Michaels, 2010). A central aspect of our knowledge of flowering time regulation is that multiple floral inductive cues are integrated into a set of flowering time integrator genes, including MADS-box genes such as *APETALA 1 (API)* and *SUPPRESSOR OF OVEREXPRESSION OF CO1 (SOC1)*, *FLOWERING LOCUS T (FT)*, and a plant-specific transcription factor *LEAFY (LFY)* (Amasino and Michaels, 2010; Lee and Lee, 2010; Srikanth and Schmid, 2011). In this Review, I begin with a brief description of the five flowering time pathways in *A. thaliana*. Then I turn to a discussion of the molecular basis of the age pathway and how age cues are integrated into other flowering inductive cues.

Flowering behaviour in *A. thaliana* can be divided into two types, winter annual and rapid cycling, based on their requirement for a prolonged exposure to low temperature, a treatment called vernalization (reviewed in Heo and Sung,

2011; Song *et al.*, 2012; Song *et al.*, 2013). Winter annual types are late flowering and such a late-flowering phenotype can be eliminated by vernalization. Genetic studies have revealed that *FLOWERING LOCUS C (FLC)*, a MADS-box gene, acts as the master regulator in vernalization pathway. Before vernalization, *FRIGIDA (FRI)*, a gene of unknown biochemical function, activates *FLC* (Choi *et al.*, 2011). *FLC* delays flowering through repressing *FT* in the leaves and *SOC1* at shoot apex (Searle *et al.*, 2006). Transcription of *FLC* rapidly decreases in response to vernalization. It has been demonstrated that the repression of *FLC* by cold is regulated by complex mechanisms involving long-non-coding RNAs (lncRNAs), histone modification and higher order chromatin assembly (Crevillen *et al.*, 2013; Rosa *et al.*, 2013; Song *et al.*, 2012; Sun *et al.*, 2013; Zografos and Sung, 2012). In contrast to winter annual, rapid-cycling accessions are early flowering in the absence of vernalization, which is often due to naturally occurring mutations in *FRI* (Johanson *et al.*, 2000).

A. thaliana is a long day plant, in which the onset of flowering is accelerated when the length of daylight is prolonged compared with darkness. Molecular and genetic analyses demonstrate that the seasonal changes in day length are measured by *CONSTANS (CO)*, which encodes a zinc finger and CCT-domain-containing transcription factor (Putterill *et al.*, 1995). *co* mutants show delayed flowering in long days but not in short days (Putterill *et al.*, 1995). Classical physiological experiments reveal that the floral transition in response to day length involves a systemic signal formed in the leaves that induces floral transition at the SAM. Consistent with this notion, *CO* is mainly expressed in leaf vascular tissues (An *et al.*, 2004). *CO* expression is regulated by light at both the transcriptional and post-transcriptional level. In short days, the expression of *CO* peaks after dusk, so that *CO* protein is subjected to COP1-mediated degradation (Jang *et al.*, 2008; Liu *et al.*, 2008; Valverde *et al.*, 2004; Yanovsky and Kay, 2002). In contrast, *CO* expression coincides with light in long days, which leads to stabilization of *CO*. The accumulation of *CO* leads to the activation of *FT*, which encodes a putative phosphatidylethanolamine-binding protein. With the help of an endoplasmic reticulum (ER) membrane protein, FT-INTERACTING PROTEIN1 (FTIP1), *FT* proteins move from the leaves to the shoot apex (Corbesier *et al.*, 2007; Jaeger and Wigge, 2007; Lin *et al.*, 2007; Liu *et al.*, 2012; Mathieu *et al.*, 2007). In the SAM, *FT*, by binding with the transcription factor *FD*, activates the expression of *LFY* and MADS-box genes, such as *API* and *SOC1*, and thereby induces flowering (Abe *et al.*, 2005; Kobayashi and Weigel, 2007; Wigge *et al.*, 2005).

In addition to photoperoid and vernalization, GAs also play a critical role in flowering time. The GA biosynthetic mutant, *gal*, never flowers under non-inductive short day conditions (Wilson *et al.*, 1992). GA signalling transduction is mediated by ubiquitin–proteasome degradation (reviewed in Harberd, 2003; Schwechheimer and Willige, 2009). By binding to GA, GIBBERELLIN INSENSITIVE DWARF1 (GID1), a nuclear-localized GA receptor,

promotes the degradation of the transcriptional repressors called DELLAs. The *Arabidopsis* genome encodes five DELLA genes, namely *REPRESSOR OF GAI-3 (RGA)*, *GA INSENSITIVE (GAI)*, *RGA-LIKE 1 (RGL1)*, *RGL2*, and *RGL3* (Murase *et al.*, 2008). The DELLA motif, which is 17 amino acid residues long and located in the amino-terminal of DELLAs, is essential for the degradation of DELLA proteins by the proteasome (Dill *et al.*, 2001). The negative role of GA on flowering time is mediated by DELLAs such as *GAI* and *RGA*. The *gai* or *rga-Δ17* mutant, which carries the deletion of the DELLA motif, is insensitive to GA-induced proteolysis and delays flowering (Dill *et al.*, 2001; Peng *et al.*, 1997). It has been shown that *GAI* represses flowering through *SOC1* because the expression of *SOC1* is induced by GA treatment but reduced in the *gai* mutant (Moon *et al.*, 2003). Recent studies identified two GATA-type transcription factors, *GNC* (GATA, NITRATE-INDUCIBLE, CARBON-METABOLISM INVOLVED) and *GNL* (GNC-LIKE) as new components in the GA pathway (Richter *et al.*, 2010). Expression analyses indicate that *GNC* and *GNL* act downstream of DELLAs and promote flowering through activating *SOC1* (Richter *et al.*, 2013).

The autonomous pathway is constituted by a group of genes that promote flowering by suppressing *FLC*. *FCA*, *FPA*, and *FLOWERING LOCUS K (FLK)* contain an RNA-binding domain (Lim *et al.*, 2004; Macknight *et al.*, 1997; Mockler *et al.*, 2004; Schomburg *et al.*, 2001), whereas *FY* and *FLOWERING LOCUS D (FLD)* encode a protein homologous to the yeast poly-adenylation factor Pfs2p and a dimethylated histone H3 at lysine 4 (H3K4me2) demethylase, respectively (He *et al.*, 2003; Simpson *et al.*, 2003). It has been shown that *FCA*, interacting with a component of the CPSF complex, targets CstF-dependent 3' processing to the proximal site on *FLC* antisense transcripts. With the help of *FPA*, *FY*, and *FLD*, this targeted processing triggers localized histone demethylase activity and results in reduced *FLC* sense transcription (Liu *et al.*, 2010; Manzano *et al.*, 2009). Because the regulatory basis of these genes is largely unknown, the biological relevance of autonomous pathway remains unclear.

In summary, the identities and actions of the components in flowering time pathways reveal that photoperiodic pathway acts as a positive regulator of flowering, whereas other pathways promote flowering through alleviating flowering repressors. The fact that flowering eventually occurs in the photoperiodic mutants indicates that there is another flower-promoting pathway that ensures plants flower under non-inductive conditions.

miR156–SPL defines the age pathway

microRNAs (miRNAs) are 21–24 nt long, small noncoding RNAs widely distributed in animals and plants (Bartel, 2009). It has been shown that plant miRNAs regulate gene expression through transcript cleavage (Llave *et al.*, 2002; Reinhart *et al.*, 2002) and translational inhibition (Brodersen *et al.*, 2008; Chen, 2004; Li *et al.*, 2013).

miR156 is one of the most evolutionally conserved miRNAs in plants. The targets of miR156 encode a family of transcription factors, called SQUAMOSA PROMOTER BINDING LIKEs (SPLs) (Cardon *et al.*, 1999; Rhoades *et al.*, 2002). In the *Arabidopsis* genome, there are 11 *SPLs* targeted by miR156. Based on the size of encoded proteins, these *SPL* genes can be divided into two major groups, represented by *SPL3* (*SPL3*, *SPL4*, and *SPL5*) and *SPL9* (*SPL2*, *SPL6*, *SPL9*, *SPL10*, *SPL11*, *SPL13*, *SPL13-like*, and *SPL15*) (Xing *et al.*, 2010). *SPL3*, *SPL4*, and *SPL5* are much smaller than the other gene products, with the DNA-binding domain making up most of the protein. In addition, the miR156-binding site is located in the 3'UTR of *SPL3* (also *SPL4* and *SPL5*) and miR156 regulates *SPL3* expression through transcript cleavage as well as translational inhibition (Gandikota *et al.*, 2007).

Expression of miR156 is temporally regulated. Mature miR156 is highly abundant in seedlings and decreases with time (Wang *et al.*, 2009a; Wu *et al.*, 2009; Wu and Poethig, 2006). This expression pattern is observed not only in *A. thaliana*, but also in other species including *Arabidopsis alpina*, *Cardamine flexuosa*, maize, poplar, rice, and tomato (Bergonzi *et al.*, 2013; Chuck *et al.*, 2007; Wang *et al.*, 2011a; Xie *et al.*, 2012; Yoshikawa *et al.*, 2013; Zhou *et al.*, 2013). The developmental decline in miR156 is partially mediated by sugars, the products of photosynthesis (Proveniers, 2013; Yang *et al.*, 2011; Yang *et al.*, 2013; Yu *et al.*, 2013). Exogenous sugar treatment results in a rapid decrease in miR156 expression. The repression of miR156 by sugar occurs at both transcriptional and post-transcriptional levels. Consistent with these findings, the *A. thaliana chlorinal* (*chl1*) mutant, which has impaired photosynthesis, accumulates higher level of miR156 than wild type. Similarly, defoliation delays juvenile-to-adult phase transition with a concomitant rise in miR156 level (Yang *et al.*, 2011).

The importance of miR156 in flowering is inferred from the observation that overexpression of miR156 delays flowering (Jung *et al.*, 2011b; Schwab *et al.*, 2005; Schwarz *et al.*, 2008; Wu and Poethig, 2006; Yamaguchi and Abe, 2012; Zhou and Wang, 2013). Notably, the effect of miR156 overexpression on flowering is much pronounced under non-inductive short day conditions, together with the fact that miR156 expression is regulated by age, indicating that miR156 acts as an endogenous flowering cue. In agreement with this finding, overexpression of *SPL3* results in an early flowering phenotype irrespective of photoperiodic length (Wang *et al.*, 2009a; Wu and Poethig, 2006; Yamaguchi *et al.*, 2009). In contrast, the effect of *SPL9* on flowering time is ambiguous (Wang *et al.*, 2009a). Despite the fact that *SPL9* overexpression lines flower nearly at the same time as wild type, the floral transition of *SPL9* overexpression lines is clearly accelerated when the flowering time is measured by the number of leaves produced when the plants start to flower. These contradictory results can be explained by the negative role of *SPL9* on leaf initiation rate (Wang *et al.*, 2008). Indeed, overexpression of miR156 under a shoot apex specific promoter delays flowering without affecting leaf initiation rate (Wang *et al.*, 2009a). Therefore, these

results suggest an antagonistic effect between growth rate and flowering time, which prevents plants from precocious flowering.

The role of miR156 in flowering seems widely conserved among angiosperms. Overexpression of miR156 caused late flowering phenotype in many species including *A. alpina*, *C. flexuosa*, maize, potato, and rice (Bergonzi *et al.*, 2013; Bhogale *et al.*, 2014; Chuck *et al.*, 2011; Eviatar-Ribak *et al.*, 2013; Xie *et al.*, 2006; Zhou *et al.*, 2013).

Integration of age and photoperiodic pathways

Genetic studies have placed the age pathway in parallel with photoperiodic pathway. Overexpression of miR156 in an *ft* background results in a severely delay in flowering (Wang *et al.*, 2009a). In the extreme case, flowering never occurs under short day conditions. These results indicate that the miR156-mediated age pathway ensures plants flower in the absence of exogenous inductive cues.

Recent efforts have provided insight into how the miR156–SPL module regulates flowering in *A. thaliana*. In the juvenile phase, the levels of miR156-targeted *SPL* genes are low because of high amount of miR156. As plants age, the amount of miR156 is decreased, resulting in an increase in miR156-targeted *SPL* level. *SPL3* and *SPL9* promote flowering in leaves and shoot apex through two distinct mechanisms (Figure 1). In the shoot apex, *SPL3* and *SPL9* induce flowering through activating MADS-box genes, including *AP1*, *LFY*, *FUL*, and *SOC1* (Wang *et al.*, 2009a; Yamaguchi *et al.*,




Fig. 1. miR156-mediated age pathway. miR156 level is high in juvenile phase but significantly reduced in adult phase. As a result, the level of miR156-targeted *SPLs* rises, which leads to activation of miR172 and thereby reduction in the levels of miR172-targeted *AP2-like* genes (*AP2*, *TOE1*, *TOE2*, *SMZ*, and *SNZ*). In the juvenile phase, miR172-targeted *AP2-like* proteins, with the help of *TPL*, repress flowering through *FT* in leaves, and flower-promoting genes (*AP1*, *FUL*, *LFY*, and *SOC1*) in the shoot apex. In the adult phase, *CO* activates *FT* expression in leaves and miR156-targeted *SPLs* induce flowering through activating the expression of flower-promoting genes in the shoot apex.

2009). Forced expression of *SPL3* or *SPL9* under the shoot apex specific promoter leads to early flowering phenotype under both long day and short day conditions (Wang *et al.*, 2009a).

In leaves, *SPL9* activates another miRNA, miR172, by direct binding to and transcriptional activation of *MIR172b* (Wu *et al.*, 2009). miR172 targets a family of AP2-like transcription factors, including *AP2*, *SCHLAFMUTZE* (*SMZ*), *SCHNARCHZAPFEN* (*SNZ*), *TARGET OF EAT1* (*TOE1*), *TOE2*, and *TOE3* (Aukerman and Sakai, 2003; Wu *et al.*, 2009). All these miR172-targeted AP2-like transcription factors act as flowering repressors. Overexpression of miR172 causes an extremely early flowering phenotype under both short days and long days (Jung *et al.*, 2007; Yant *et al.*, 2010), whereas the increased level of *TOE1* results in late flowering (Mathieu *et al.*, 2009). Chromatin immunoprecipitation sequencing (ChIP-SEQ) analyses reveal that *TOE1* and *AP2* not only inhibit *FT* expression in leaves, but also repress many other flowering time regulators acting downstream of *FT* in the shoot apex (Mathieu *et al.*, 2009). The repression of these genes by *TOE1* is mediated by *TOPLESS* (*TPL*), a transcriptional co-repressor (Causier *et al.*, 2012; Long *et al.*, 2006). Intriguingly, *AP2* also negatively regulates miR172 and positively regulates miR156, suggesting a miR156–miR172 feedback loop in fine-tuning the flowering response (Yant *et al.*, 2010).

Previous studies have suggested that miR156-targeted *SPL* genes act downstream of photoperiodic pathway because up-regulation of *SPL3* and *SPL9* is readily detectable within 3 days after transfer of vegetative plants from short days to inductive long days, and this induction is much reduced in *co* or *ft* mutants (Schmid *et al.*, 2003). In agreement with this finding, *SPL3* has been shown to be directly regulated by *SOC1* (Jung *et al.*, 2011a) and *SPL4* expression is reduced in a *soc1 ful* double mutant (Torti *et al.*, 2012) (Figure 2). Furthermore, mutations in two *BELL1*-like homeobox genes, *PENNYWISE* (*PNY*) and *POUND-FOOLISH* (*PNF*), impair the photoperiodic induction of *SPL3*, *SPL4*, and *SPL5* (Lal *et al.*, 2011).

Another layer of crosstalk between age and photoperiodic pathways comes from the regulation of miR172 by photoperiodic length in leaves (Figure 2). *GIGANTEA* (*GI*) positively regulates *CO* transcription (Fowler *et al.*, 1999; Jung *et al.*, 2007; Park *et al.*, 1999). miR172 abundance is substantially reduced in a *gi* mutant (Jung *et al.*, 2007). It is suggested that *GI* regulates miR172 at the miRNA processing level because the level of primary transcript of *MIR172* is not accordingly reduced but elevated in a *gi* mutant.

Crosstalk between age and gibberellin pathways

Under non-inductive short day conditions, age and GA pathways play the predominant roles in flowering. *DELLA* represses flowering in both leaves and the shoot apex. Forced expression of GA-insensitive *RGA* or GA catabolic




Fig. 2. Integration of age and photoperiodic pathways. The integration of age and photoperiodic pathways take place at two levels (purple arrow lines). First, miR172 abundance is regulated by photoperiod via *GI*-mediated miRNA processing. Second, miR156-targeted *SPLs* acts not only in parallel with, but also downstream of photoperiodic pathway. The level of miR156-targeted *SPLs* is rapidly induced when *A. thaliana* plants are shifted from short days to long days. This action seems to be mediated by two *MADS*-box genes, *SOC1* and *FUL*.

genes under leaf or shoot apex specific promoters results in a late flowering phenotype (Galvao *et al.*, 2012; Porri *et al.*, 2012; Yu *et al.*, 2012). Interestingly, GA treatment does not markedly accelerate flowering in an miR156 overexpression line, indicating that GA promotes flowering partially through the miR156–*SPL* module. In light of this finding, Yu *et al.* (2012) revealed that GA and age pathways are integrated through a physical interaction between *DELLAs* (*RGA*, *GAI*, *RGL1*, *RGA2* and *RGL3*) and miR156-targeted *SPL9*-like proteins (*SPL2*, *SPL9*, *SPL10* and *SPL11*). The binding of *RGA* to *SPL9* interferes with *SPL9* transcriptional activities on *MIR172b*, *SOC1*, and *FUL*. As a result, *DELLA* delays flowering by reducing *FT* expression through repressing miR172 in leaves, whereas it inhibits floral transition by repressing *SOC1* and *FUL* in the shoot apex (Figure 3).

Integration of age and vernalization pathways

In *A. thaliana*, plants become competent to vernalization after germination. However, recent studies indicate that age regulates the timing of sensitivity in response to vernalization in *A. alpina* and *C. flexuosa*, two polycarpic perennials closely related to *A. thaliana*. Independently, Wang *et al.* and Zhou *et al.* revealed that young *A. alpina* and *C. flexuosa* are insensitive to cold treatment (Wang *et al.*, 2011b; Zhou *et al.*, 2013). This flowering behaviour is mediated by the levels of miR156 and miR172 (Bergonzi *et al.*, 2013; Zhou *et al.*, 2013). Overexpression of miR156 prevents flowering in response to




Fig. 3. Integration of Age and GA pathways. (A) DELLA acts as a repressor of miR156-targeted SPL. The binding of DELLA with SPL inhibits the transcriptional activation of SPL targets, such as miR172, *FUL*, and *SOC1*. (B) DELLA represses flowering in two distinct mechanisms. The binding of DELLA with miR156-targeted SPLs compromises the activation of miR172 in leaves, and the activation of MADS-box genes (*SOC1* and *FUL*) in the shoot apex. In addition, it has been shown that DELLAs indirectly repress *SOC1* through two GATA transcription factors, GNC and GNL.

vernalization, whereas the reduced activity of miR156 or *PERPETUAL FLOWERING2* (*PEP2*, an miR172-targeted *AP2-like* gene) results in an accelerated acquisition of floral competence in response to vernalization. In addition, *A. alpina* *TERMINAL FLOWER1* (*AaTFL1*) was found to block flowering of young *A. alpina* plants exposed to vernalization (Wang *et al.*, 2011b). The integration of age and vernalization pathways thus offers an advantage for the perennial growth habit by ensuring that plants do not flower until they develop axillary vegetative shoots and sufficient biomass.

Although the role of miR156 and miR172 in setting a threshold for the sensitivity in response to vernalization is conserved between *A. alpina* and *C. flexuosa*, the underlying molecular mechanism differs in the following two aspects (Fig. 4). First, *C. flexuosa* *FLC* expression is not reduced when miR172-targeted *AP2* group genes are suppressed by miR172 overexpression, whereas *PEP1*, the *A. alpina* *FLC* orthologue (Wang *et al.*, 2009b), is decreased in the *pep2* mutant (Fig. 4). Second, the expression of flowering activator miR172 is coupled with the flowering repressor miR156 in *A. thaliana*, maize, rice, and poplar. In *C. flexuosa*, miR172 is similarly linked to miR156, whereas it seems that *A. alpina* is an exception from this rule (Fig. 4). Interestingly, although the level of miR172 is not increased during vegetative phase, a rise in miR172 abundance is observed in developing floral primordia, which leads to alleviate the flowering repressive effect of miR172-targeted *AP2-like* proteins (Bergonzi *et al.*, 2013).

The above two differences reflect different strategies in the two perennial species. In *A. alpina*, because *PEP2* positively regulates *FLC*, miR172-targeted *AP2-like* genes have to be uncoupled from miR156 and its SPL targets. Otherwise, the age-dependent increase in miR172 will cause a loss of *FLC* activity and thus promote flowering. Conversely, in *C. flexuosa*, because miR172 has remained under the control of miR156–SPL module, *FLC* has to be uncoupled from miR172-targeted *AP2-like* genes (Fig. 4).

Future directions

The past 20 years have witnessed a great increase in our knowledge of the basic molecular mechanisms of flowering. Most remarkably, functional genetic studies in *A. thaliana* and rice have identified signalling pathways that act as master regulators of floral transition and that are conserved in monocots and dicots. Growing evidence suggests that the integration of each floral inductive cue varies in different species. As described above, although both vernalization and age pathways operate in *A. thaliana*, this species does not have a pronounced age-dependent vernalization response. Thus, a major challenge in the future will be to understand how the flowering pathways are differentially regulated and integrated in different species.

The identification of sugar as an upstream regulator of miR156 suggests that sugar may play an important role in flowering. Consistently, trehalose-6-phosphate (T6P), a disaccharide molecule, was recently revealed as a new




Fig. 4. Integration of age and vernalization pathways. (A) In *A. alpina*, miR172 is not regulated by an miR156-mediated age pathway. PEP2 (an *A. alpina* AP2) activates the expression of PEP1, an *FLC* orthologue in *A. alpina* (red arrow line). Note that miR172 level is not changed with age but arises in developing floral primordia. The dashed arrow line indicates an indirect activation of SOC1 by miR156-targeted AaSPL. (B) In *C. flexuosa*, the miR156–SPL module is directly connected to the miR172–AP2 module (red arrow line). The expression of CfFLC is not regulated by CfTOE1 (a *C. flexuosa* miR172-targeted AP2-like gene).

regulator of flowering (van Dijken *et al.*, 2004; Wahl *et al.*, 2013). Therefore, another challenge in future is to explore the means by which carbohydrate or energetic status regulates flowering.

Acknowledgements

The work in the Wang lab is supported by the grants from National Natural Science Foundation of China (31222029; 91217306), State Key Basic Research Program of China (2013CB127000), Shanghai Pujiang Program (12PJ1409900), Recruitment Program of Global Experts (China), and Key Research Program grant from NKLPMPG (SIPPE, SIBS). I thank Yan Qian, Hongtao Liu, and members in Wang lab for helpful comments on the manuscript and apologize to those authors whose work is not cited owing to space limitations.

References

Abe M, Kobayashi Y, Yamamoto S, Daimon Y, Yamaguchi A, Ikeda Y, Ichinoki H, Notaguchi M, Goto K, Araki T. 2005. FD, a bZIP protein mediating signals from the floral pathway integrator FT at the shoot apex. *Science* **309**, 1052–1056.

Amasino RM, Michaels SD. 2010. The timing of flowering. *Plant Physiology* **154**, 516–520.

An H, Roussot C, Suarez-Lopez P *et al.* 2004. CONSTANS acts in the phloem to regulate a systemic signal that induces photoperiodic flowering of *Arabidopsis*. *Development* **131**, 3615–3626.

Andres F, Coupland G. 2012. The genetic basis of flowering responses to seasonal cues. *Nature Reviews Genetics* **13**, 627–639.

Aukerman MJ, Sakai H. 2003. Regulation of flowering time and floral organ identity by a MicroRNA and its *APETALA2*-like target genes. *Plant Cell* **15**, 2730–2741.

Bäurle I, Dean C. 2006. The timing of developmental transitions in plants. *Cell* **125**, 655–664.

Bartel DP. 2009. MicroRNAs: target recognition and regulatory functions. *Cell* **136**, 215–233.

Bergonzi S, Albani MC, Ver Loren van Themaat E, Nordstrom KJ, Wang R, Schneeberger K, Moerland PD, Coupland G. 2013. Mechanisms of age-dependent response to winter temperature in perennial flowering of *Arabidopsis alpina*. *Science* **340**, 1094–1097.

Bhogale S, Mahajan AS, Natarajan B, Rajabhoj M, Thulasiram HV, Banerjee AK. 2014. MicroRNA156: A potential graft-transmissible microRNA that modulates plant architecture and tuberization in *Solanum tuberosum* ssp. *andigena*. *Plant Physiology* **164**, 1011–1027.

Brodersen P, Sakvarelidze-Achard L, Bruun-Rasmussen M, Dunoyer P, Yamamoto YY, Sieburth L, Voinnet O. 2008. Widespread translational inhibition by plant miRNAs and siRNAs. *Science* **320**, 1185–1190.

Cardon G, Hohmann S, Klein J, Nettlesheim K, Saedler H, Huijser P. 1999. Molecular characterisation of the *Arabidopsis* SBP-box genes. *Gene* **237**, 91–104.

Causier B, Ashworth M, Guo W, Davies B. 2012. The TOPLESS interactome: a framework for gene repression in *Arabidopsis*. *Plant Physiology* **158**, 423–438.

Chen X. 2004. A microRNA as a translational repressor of *APETALA2* in *Arabidopsis* flower development. *Science* **303**, 2022–2025.

Choi K, Kim J, Hwang HJ, Kim S, Park C, Kim SY, Lee I. 2011. The FRIGIDA complex activates transcription of *FLC*, a strong flowering

repressor in *Arabidopsis*, by recruiting chromatin modification factors. *Plant Cell* **23**, 289–303.

Chuck G, Cigan AM, Saeteurn K, Hake S. 2007. The heterochronic maize mutant *Corngrass1* results from overexpression of a tandem microRNA. *Nature Genetics* **39**, 544–549.

Chuck GS, Tobias C, Sun L, Kraemer F, Li C, Dibble D, Arora R, Bragg JN, Vogel JP, Singh S, Simmons BA, Pauly M, Hake S. 2011. Overexpression of the maize *Corngrass1* microRNA prevents flowering, improves digestibility, and increases starch content of switchgrass. *Proceedings of the National Academy of Sciences, USA* **108**, 17550–17555.

Corbesier L, Vincent C, Jang S et al. 2007. FT protein movement contributes to long-distance signaling in floral induction of *Arabidopsis*. *Science* **316**, 1030–1033.

Crevillen P, Sonmez C, Wu Z, Dean C. 2013. A gene loop containing the floral repressor *FLC* is disrupted in the early phase of vernalization. *The EMBO Journal* **32**, 140–148.

Dill A, Jung HS, Sun TP. 2001. The DELLA motif is essential for gibberellin-induced degradation of RGA. *Proceedings of the National Academy of Sciences, USA* **98**, 14162–14167.

Eviatar-Ribak T, Shalit-Kaneh A, Chappell-Maor L, Amsellem Z, Eshed Y, Lifschitz E. 2013. A cytokinin-activating enzyme promotes tuber formation in tomato. *Current Biology* **23**, 1057–1064.

Fowler S, Lee K, Onouchi H, Samach A, Richardson K, Morris B, Coupland G, Putterill J. 1999. *GIGANTEA*: a circadian clock-controlled gene that regulates photoperiodic flowering in *Arabidopsis* and encodes a protein with several possible membrane-spanning domains. *The EMBO Journal* **18**, 4679–4688.

Galvao VC, Horrer D, Kuttner F, Schmid M. 2012. Spatial control of flowering by DELLA proteins in *Arabidopsis thaliana*. *Development* **139**, 4072–4082.

Gandikota M, Birkenbihl RP, Hohmann S, Cardon GH, Saedler H, Huijser P. 2007. The miRNA156/157 recognition element in the 3' UTR of the *Arabidopsis* SBP box gene *SPL3* prevents early flowering by translational inhibition in seedlings. *The Plant Journal* **49**, 683–693.

Harberd NP. 2003. Botany. Relieving DELLA restraint. *Science* **299**, 1853–1854.

He Y, Michaels SD, Amasino RM. 2003. Regulation of flowering time by histone acetylation in *Arabidopsis*. *Science* **302**, 1751–1754.

Heo JB, Sung S. 2011. Vernalization-mediated epigenetic silencing by a long intronic noncoding RNA. *Science* **331**, 76–79.

Huijser P, Schmid M. 2011. The control of developmental phase transitions in plants. *Development* **138**, 4117–4129.

Jaeger KE, Wigge PA. 2007. FT protein acts as a long-range signal in *Arabidopsis*. *Current Biology* **17**, 1050–1054.

Jang S, Marchal V, Panigrahi KC, Wenkel S, Soppe W, Deng XW, Valverde F, Coupland G. 2008. *Arabidopsis* COP1 shapes the temporal pattern of CO accumulation conferring a photoperiodic flowering response. *The EMBO Journal* **27**, 1277–1288.

Johanson U, West J, Lister C, Michaels S, Amasino R, Dean C. 2000. Molecular analysis of *FRIGIDA*, a major determinant of natural variation in *Arabidopsis* flowering time. *Science* **290**, 344–347.

Jung JH, Ju Y, Seo PJ, Lee JH, Park CM. 2011a. The SOC1–SPL module integrates photoperiod and gibberellic acid signals to control flowering time in *Arabidopsis*. *The Plant Journal* **69**, 577–588.

Jung JH, Seo PJ, Kang SK, Park CM. 2011b. miR172 signals are incorporated into the miR156 signaling pathway at the *SPL3/4/5* genes in *Arabidopsis* developmental transitions. *Plant Molecular Biology* **76**, 35–45.

Jung JH, Seo YH, Seo PJ, Reyes JL, Yun J, Chua NH, Park CM. 2007. The *GIGANTEA*-regulated microRNA172 mediates photoperiodic flowering independent of *CONSTANS* in *Arabidopsis*. *Plant Cell* **19**, 2736–2748.

Kobayashi Y, Weigel D. 2007. Move on up, it's time for change—mobile signals controlling photoperiod-dependent flowering. *Genes and Development* **21**, 2371–2384.

Lal S, Pacis LB, Smith HM. 2011. Regulation of the *SQUAMOSA PROMOTER-BINDING PROTEIN-LIKE* genes/microRNA156 module

by the homeodomain proteins PENNYWISE and POUND-FOOLISH in *Arabidopsis*. *Molecular Plant* **4**, 1123–1132.

Lee J, Lee I. 2010. Regulation and function of *SOC1*, a flowering pathway integrator. *Journal of Experimental Botany* **61**, 2247–2254.

Li S, Liu L, Zhuang X et al. 2013. MicroRNAs inhibit the translation of target mRNAs on the endoplasmic reticulum in *Arabidopsis*. *Cell* **153**, 562–574.

Lim MH, Kim J, Kim YS, Chung KS, Seo YH, Lee I, Kim J, Hong CB, Kim HJ, Park CM. 2004. A new *Arabidopsis* gene, *FLK*, encodes an RNA binding protein with K homology motifs and regulates flowering time via FLOWERING LOCUS C. *Plant Cell* **16**, 731–740.

Lin MK, Belanger H, Lee YJ et al. 2007. FLOWERING LOCUS T protein may act as the long-distance florigenic signal in the cucurbits. *Plant Cell* **19**, 1488–1506.

Liu F, Marquardt S, Lister C, Swiezewski S, Dean C. 2010. Targeted 3' processing of antisense transcripts triggers *Arabidopsis* *FLC* chromatin silencing. *Science* **327**, 94–97.

Liu L, Liu C, Hou X, Xi W, Shen L, Tao Z, Wang Y, Yu H. 2012. FTIP1 is an essential regulator required for florigen transport. *PLoS Biology* **10**, e1001313.

Liu LJ, Zhang YC, Li QH, Sang Y, Mao J, Lian HL, Wang L, Yang HQ. 2008. COP1-mediated ubiquitination of *CONSTANS* is implicated in cryptochrome regulation of flowering in *Arabidopsis*. *Plant Cell* **20**, 292–306.

Llave C, Xie Z, Kasschau KD, Carrington JC. 2002. Cleavage of Scarecrow-like mRNA targets directed by a class of *Arabidopsis* miRNA. *Science* **297**, 2053–2056.

Long JA, Ohno C, Smith ZR, Meyerowitz EM. 2006. TOPLESS regulates apical embryonic fate in *Arabidopsis*. *Science* **312**, 1520–1523.

Macknight R, Bancroft I, Page T et al. 1997. *FCA*, a gene controlling flowering time in *Arabidopsis*, encodes a protein containing RNA-binding domains. *Cell* **89**, 737–745.

Manzano D, Marquardt S, Jones AM, Baurle I, Liu F, Dean C. 2009. Altered interactions within FY/AtCPSF complexes required for *Arabidopsis* *FCA*-mediated chromatin silencing. *Proceedings of the National Academy of Sciences, USA* **106**, 8772–8777.

Mathieu J, Warthmann N, Kuttner F, Schmid M. 2007. Export of FT protein from phloem companion cells is sufficient for floral induction in *Arabidopsis*. *Current Biology* **17**, 1055–1060.

Mathieu J, Yant LJ, Murdter F, Kuttner F, Schmid M. 2009. Repression of flowering by the miR172 target *SMZ*. *PLoS Biology* **7**, e1000148.

Mockler TC, Yu X, Shalitin D et al. 2004. Regulation of flowering time in *Arabidopsis* by K homology domain proteins. *Proceedings of the National Academy of Sciences, USA* **101**, 12759–12764.

Moon J, Suh SS, Lee H, Choi KR, Hong CB, Paek NC, Kim SG, Lee I. 2003. The *SOC1* MADS-box gene integrates vernalization and gibberellin signals for flowering in *Arabidopsis*. *The Plant Journal* **35**, 613–623.

Murase K, Hirano Y, Sun TP, Hakoshima T. 2008. Gibberellin-induced DELLA recognition by the gibberellin receptor GID1. *Nature* **456**, 459–463.

Park DH, Somers DE, Kim YS, Choy YH, Lim HK, Soh MS, Kim HJ, Kay SA, Nam HG. 1999. Control of circadian rhythms and photoperiodic flowering by the *Arabidopsis* *GIGANTEA* gene. *Science* **285**, 1579–1582.

Peng J, Carol P, Richards DE, King KE, Cowling RJ, Murphy GP, Harberd NP. 1997. The *Arabidopsis* *GAI* gene defines a signaling pathway that negatively regulates gibberellin responses. *Genes and Development* **11**, 3194–3205.

Poethig RS. 2003. Phase change and the regulation of developmental timing in plants. *Science* **301**, 334–336.

Poethig RS. 2013. Vegetative phase change and shoot maturation in plants. *Current Topics in Developmental Biology* **105**, 125–152.

Porri A, Torti S, Romera-Branchat M, Coupland G. 2012. Spatially distinct regulatory roles for gibberellins in the promotion of flowering of *Arabidopsis* under long photoperiods. *Development* **139**, 2198–2209.

Proveniers M. 2013. Sugars speed up the circle of life. *elife* **2**, e00625.

Putterill J, Robson F, Lee K, Simon R, Coupland G. 1995. The *CONSTANS* gene of *Arabidopsis* promotes flowering and encodes a

- protein showing similarities to zinc finger transcription factors. *Cell* **80**, 847–857.
- Reinhart BJ, Weinstein EG, Rhoades MW, Bartel B, Bartel DP.** 2002. MicroRNAs in plants. *Genes and Development* **16**, 1616–1626.
- Rhoades MW, Reinhart BJ, Lim LP, Burge CB, Bartel B, Bartel DP.** 2002. Prediction of plant microRNA targets. *Cell* **110**, 513–520.
- Richter R, Bastakis E, Schwechheimer C.** 2013. Cross-repressive interactions between SOC1 and the GATAs GNC and GNL/CGA1 in the control of greening, cold tolerance, and flowering time in *Arabidopsis*. *Plant Physiology* **162**, 1992–2004.
- Richter R, Behringer C, Muller IK, Schwechheimer C.** 2010. The GATA-type transcription factors GNC and GNL/CGA1 repress gibberellin signaling downstream from DELLA proteins and PHYTOCHROME-INTERACTING FACTORS. *Genes and Development* **24**, 2093–2104.
- Rosa S, De Lucia F, Mylne JS, Zhu D, Ohmido N, Pendle A, Kato N, Shaw P, Dean C.** 2013. Physical clustering of *FLC* alleles during Polycomb-mediated epigenetic silencing in vernalization. *Genes and Development* **27**, 1845–1850.
- Schmid M, Uhlenhaut NH, Godard F, Demar M, Bressan R, Weigel D, Lohmann JU.** 2003. Dissection of floral induction pathways using global expression analysis. *Development* **130**, 6001–6012.
- Schomburg FM, Patton DA, Meinke DW, Amasino RM.** 2001. *FPA*, a gene involved in floral induction in *Arabidopsis*, encodes a protein containing RNA-recognition motifs. *Plant Cell* **13**, 1427–1436.
- Schwab R, Palatnik JF, Rieger M, Schommer C, Schmid M, Weigel D.** 2005. Specific effects of microRNAs on the plant transcriptome. *Developmental Cell* **8**, 517–527.
- Schwarz S, Grande AV, Bujdoso N, Saedler H, Huijser P.** 2008. The microRNA regulated SBP-box genes *SPL9* and *SPL15* control shoot maturation in *Arabidopsis*. *Plant Molecular Biology* **67**, 183–195.
- Schwechheimer C, Willige BC.** 2009. Shedding light on gibberellin acid signalling. *Current Opinion in Plant Biology* **12**, 57–62.
- Searle I, He Y, Turck F, Vincent C, Fornara F, Krober S, Amasino RA, Coupland G.** 2006. The transcription factor *FLC* confers a flowering response to vernalization by repressing meristem competence and systemic signaling in *Arabidopsis*. *Genes and Development* **20**, 898–912.
- Simpson GG, Dijkwel PP, Quesada V, Henderson I, Dean C.** 2003. *FY* is an RNA 3' end-processing factor that interacts with *FCA* to control the *Arabidopsis* floral transition. *Cell* **113**, 777–787.
- Song J, Angel A, Howard M, Dean C.** 2012. Vernalization—a cold-induced epigenetic switch. *Journal of Cell Science* **125**, 3723–3731.
- Song J, Irwin J, Dean C.** 2013. Remembering the prolonged cold of winter. *Current Biology* **23**, R807–811.
- Srikanth A, Schmid M.** 2011. Regulation of flowering time: all roads lead to Rome. *Cellular and Molecular Life Sciences* **68**, 2013–2037.
- Sun Q, Csorba T, Skourti-Stathaki K, Proudfoot NJ, Dean C.** 2013. R-loop stabilization represses antisense transcription at the *Arabidopsis FLC* locus. *Science* **340**, 619–621.
- Torti S, Fornara F, Vincent C, Andres F, Nordstrom K, Gobel U, Knoll D, Schoof H, Coupland G.** 2012. Analysis of the *Arabidopsis* shoot meristem transcriptome during floral transition identifies distinct regulatory patterns and a leucine-rich repeat protein that promotes flowering. *Plant Cell* **24**, 444–462.
- Valverde F, Mouradov A, Soppe W, Ravenscroft D, Samach A, Coupland G.** 2004. Photoreceptor regulation of *CONSTANS* protein in photoperiodic flowering. *Science* **303**, 1003–1006.
- van Dijken AJ, Schluempmann H, Smeekens SC.** 2004. *Arabidopsis* trehalose-6-phosphate synthase 1 is essential for normal vegetative growth and transition to flowering. *Plant Physiology* **135**, 969–977.
- Wahl V, Ponnu J, Schlereth A, Arrivault S, Langenecker T, Franke A, Feil R, Lunn JE, Stitt M, Schmid M.** 2013. Regulation of flowering by trehalose-6-phosphate signaling in *Arabidopsis thaliana*. *Science* **339**, 704–707.
- Wang JW, Czech B, Weigel D.** 2009a. miR156-regulated SPL transcription factors define an endogenous flowering pathway in *Arabidopsis thaliana*. *Cell* **138**, 738–749.
- Wang JW, Park MY, Wang LJ, Koo Y, Chen XY, Weigel D, Poethig RS.** 2011a. miRNA control of vegetative phase change in trees. *PLoS Genetics* **7**, e1002012.
- Wang JW, Schwab R, Czech B, Mica E, Weigel D.** 2008. Dual effects of miR156-targeted *SPL* genes and *CYP78A5/KLUH* on plastochron length and organ size in *Arabidopsis thaliana*. *Plant Cell* **20**, 1231–1243.
- Wang R, Albani MC, Vincent C, Bergonzi S, Luan M, Bai Y, Kiefer C, Castillo R, Coupland G.** 2011b. *Aa TFL1* confers an age-dependent response to vernalization in perennial *Arabis alpina*. *Plant Cell* **23**, 1307–1321.
- Wang R, Farrona S, Vincent C, Joecker A, Schoof H, Turck F, Alonso-Blanco C, Coupland G, Albani MC.** 2009b. *PEP1* regulates perennial flowering in *Arabis alpina*. *Nature* **459**, 423–427.
- Wigge PA, Kim MC, Jaeger KE, Busch W, Schmid M, Lohmann JU, Weigel D.** 2005. Integration of spatial and temporal information during floral induction in *Arabidopsis*. *Science* **309**, 1056–1059.
- Wilson RN, Heckman JW, Somerville CR.** 1992. Gibberellin is required for flowering in *Arabidopsis thaliana* under short days. *Plant Physiology* **100**, 403–408.
- Wu G, Park MY, Conway SR, Wang JW, Weigel D, Poethig RS.** 2009. The sequential action of miR156 and miR172 regulates developmental timing in *Arabidopsis*. *Cell* **138**, 750–759.
- Wu G, Poethig RS.** 2006. Temporal regulation of shoot development in *Arabidopsis thaliana* by miR156 and its target *SPL3*. *Development* **133**, 3539–3547.
- Xie K, Shen J, Hou X, Yao J, Li X, Xiao J, Xiong L.** 2012. Gradual increase of miR156 regulates temporal expression changes of numerous genes during leaf development in rice. *Plant Physiology* **158**, 1382–1394.
- Xie K, Wu C, Xiong L.** 2006. Genomic organization, differential expression, and interaction of *SQUAMOSA* promoter-binding-like transcription factors and microRNA156 in rice. *Plant Physiology* **142**, 280–293.
- Xing S, Salinas M, Hohmann S, Berndtgen R, Huijser P.** 2010. miR156-targeted and nontargeted SBP-box transcription factors act in concert to secure male fertility in *Arabidopsis*. *Plant Cell* **22**, 3935–3950.
- Yamaguchi A, Abe M.** 2012. Regulation of reproductive development by non-coding RNA in *Arabidopsis*: to flower or not to flower. *Journal of Plant Research* **125**, 693–704.
- Yamaguchi A, Wu MF, Yang L, Wu G, Poethig RS, Wagner D.** 2009. The microRNA-regulated SBP-Box transcription factor *SPL3* is a direct upstream activator of *LEAFY*, *FRUITFULL*, and *APETALA1*. *Developmental Cell* **17**, 268–278.
- Yang L, Conway SR, Poethig RS.** 2011. Vegetative phase change is mediated by a leaf-derived signal that represses the transcription of miR156. *Development* **138**, 245–249.
- Yang L, Xu M, Koo Y, He J, Poethig RS.** 2013. Sugar promotes vegetative phase change in *Arabidopsis thaliana* by repressing the expression of *MIR156A* and *MIR156C*. *elife* **2**, e00260.
- Yanovsky MJ, Kay SA.** 2002. Molecular basis of seasonal time measurement in *Arabidopsis*. *Nature* **419**, 308–312.
- Yant L, Mathieu J, Dinh TT, Ott F, Lanz C, Wollmann H, Chen X, Schmid M.** 2010. Orchestration of the floral transition and floral development in *Arabidopsis* by the bifunctional transcription factor *APETALA2*. *Plant Cell* **22**, 2156–2170.
- Yoshikawa T, Ozawa S, Sentoku N, Itoh J, Nagato Y, Yokoi S.** 2013. Change of shoot architecture during juvenile-to-adult phase transition in soybean. *Planta* **238**, 229–237.
- Yu S, Cao L, Zhou CM, Zhang TQ, Lian H, Sun Y, Wu JQ, Huang JR, Wang GD, Wang JW.** 2013. Sugar is an endogenous cue for juvenile-to-adult phase transition in plants. *elife* **2**, e00269.
- Yu S, Galvao VC, Zhang YC, Horrer D, Zhang TQ, Hao YH, Feng YQ, Wang S, Schmid M, Wang JW.** 2012. Gibberellin regulates the *Arabidopsis* floral transition through miR156-targeted *SQUAMOSA* PROMOTER BINDING-LIKE transcription factors. *Plant Cell* **24**, 3320–3332.
- Zhou CM, Wang JW.** 2013. Regulation of flowering time by microRNAs. *Journal of Genetics and Genomics* **40**, 211–215.
- Zhou CM, Zhang TQ, Wang X, Yu S, Lian H, Tang H, Feng ZY, Zoumova-Lihova J, Wang JW.** 2013. Molecular basis of age-dependent vernalization in *Cardamine flexuosa*. *Science* **340**, 1097–1100.
- Zografos BR, Sung S.** 2012. Vernalization-mediated chromatin changes. *Journal of Experimental Botany* **63**, 4343–4348.